Brain Research, 559 (1991) 358–359
Elsevier

Author Index

Andrews, B.J., see Granat, M.H., 344
Aoki, E., Semba, R. and Kashiwamata, S., Evidence for the presence of L-arginine in the glial components of the peripheral nervous system, 159
Appenteng, K., see Saha, S., 145
Ashe, J.H., see Metherate, R., 163

Bain, J.R., see Evans, P.J., 315
Barnstable, C.J., see Naegele, J.R., 118
Batten, T.F.C., see Saha, S., 145
Baudry, M., see Tocco, G., 168
Baudry, M., see Tocco, G., 331
Baxendale, R.H., see Granat, M.H., 344
Bayer, V.E. and Pickel, V.M., GABA-labeled terminals form proportionally more synapses with dopaminergic neurons containing low densities of tyrosine hydroxylase-immunoreactivity in rat ventral tegmental area, 44
Berecek, K.H., see Swords, B.H., 10
Besharse, J.C., see Grace, M.S., 56
Besson, M.J., see Le Boulch, N.L., 200
Betancur, C., see Neveu, P.J., 347
Biella, G., Panara, C., Pecile, A. and Sotgiu, M.L., Facilitatory role of calcitonin gene-related peptide (CGRP) on excitation induced by substance P (SP) and noxious stimuli in rat spinal dorsal horn neurons. An iontophoretic study in vivo, 352
Blaustein, J.D., see Delville, Y., 191
Blomqvist, L., see Sjögren, A., 276
Boldry, R.C., Willins, D.L., Wallace, L.J. and Uretsky, N.J., The role of endogenous dopamine in the hypermotility response to intra-accumbens AMPA, 100
Boyes, B.E., see Ousky, J.R., 322
Brammer, G.L., see Raleigh, M.J., 181
Brunello, N., see Reynolds, M.E., 149

Cahill, G.M., see Grace, M.S., 56
Connelly, C.D., see Raffa, R.B., 211
Crain, S.M., see Shen, K.-F., 130
Creese, I., see Tepper, J.M., 283
Cunningham, S.T., Steinman, J.L., Whipple, B., Mayer, A.D. and Komisaruk, B.R., Differential roles of hypogastric and pelvic nerves in the analgesic and motoric effects of vaginocervical stimulation in rats, 337

DeGeorge, J.J., see Nariai, T., 1
Delville, Y. and Blaustein, J.D., A site for estradiol priming of progesterone-facilitated sexual receptivity in the ventrolateral hypothalamus of female guinea pigs, 191
Devgan, K.K., see Tocco, G., 331
Dragunow, M., see Jansen, K.L.R., 172
Duffy, P., see Samathanam, G.K., 37

Evans, P.J., Bain, J.R., Mackinnon, S.E., Makino, A.P. and Hunter, D.A., Selective reinnervation: a comparison of recovery following microsuture and conduit nerve repair, 315

Faull, R.L.M., see Jansen, K.L.R., 172

Gábriel, R., Zhu, B. and Straznicky, C., Tyrosine hydroxylase-immunoreactive elements in the distal retina of *Bufo marinus*: a light and electron microscopic study, 225
Gauchy, C., see Le Boulch, N.L., 200
Grace, M.S., Cahill, G.M. and Besharse, J.C., Melatonin deacetylation: retinal vertebrate class distribution and *Xenopus laevis* tissue distribution, 56
Granat, M.H., Nicol, D.J., Baxendale, R.H. and Andrews, B.J., Dishabituation of the flexion reflex in spinal cord-injured man and its applicatioin in the restoration of gait, 344
Grant, S.J. and Highfield, D.A., Extracellular characteristics of putative cholinergic neurons in the rat laterodorsal tegmental nucleus, 64
Gunnet, J.W., see Lookingland, K.J., 82

Hama, H., see Yamaguchi, K., 261
Hamada, M., see Ueyama, T., 293
Hano, T., see Ueyama, T., 293
Hauge, S.A., see Tocco, G., 331
Heiligenberg, W., see Zupanc, G.K.H., 249
Highfield, D.A., see Grant, S.J., 64
Ho, L.T., see Zochodne, D.W., 89
Hollingworth, R.M., see Raffa, R.B., 211
Hori, T., see Tanaka, T., 154
Hunter, .A., see Evans, P.J., 315

Ireland, L.M., see Lookingland, K.J., 82

Jansen, K.L.R., Faull, R.L.M., Dragunow, M. and Leslie, R.A., Autoradiographic distribution of sigma receptors in human neocortex, hippocampus, basal ganglia, cerebellum, pineal and pituitary glands, 172

Kalivas, P.W., see Samathanam, G.K., 37
Kalivas, P.W., see Sorg, B.A., 29
Karius, D.R., Ling, L. and Speck, D.F., Lesions of the rostral dorsolateral pons have no effect on afferent-evoked inhibition of inspiration, 22
Kashiwamata, S., see Aoki, E., 159
Kayahara, T., see Yasui, Y., 139
Kendig, J.J., see Woodley, S.J., 17
Komisaruk, .R., see Cunningham, S.T., 337
Kondo, S., see Tanaka, T., 154
Korn, H., see Petrov, T., 75
Ku, Y.-H. and Wang, X.-Q., Favored patterns in spontaneous spike trains, 241

Lamour, Y., see Nariai, T., 1
Le Boulch, N.L., Truong-Ngoc, N.A., Gauchy, C. and Besson, M.J., In vivo release of newly synthesized [^{3}H]GABA in the substantia nigra of the rat: relative contribution of GABA striato-pallidonigral afferents and nigral GABA neurons, 200
Le Moal, M., see Neveu, P.J., 347
Ledeen, R.W., see Shen, K.-F., 130
Leslie, R.A., see Jansen, K.L.R., 172
Li, J., see Luo, P., 267
Lichtman, A.H. and Martin, B.R., Cannabinoid-induced antinociception is mediated by a spinal α_2-noradrenergic mechanism, 309
Lindå, H., see Sjögren, A., 276
Ling, L., see Karius, D.R., 22
Long, J.P., see Ma, S., 297
Lookingland, K.J., Ireland, L.M., Gunnet, J.W., Manzanares, J., Tian, Y. and Moore, K.E., 3-Methoxy-4-hydroxyphenylethyleneglycol concentrations in discrete hypothalamic nuclei reflect the activity of noradrenergic neurons, 82
Lu, C.-L., see Shaikh, M.B., 109
Luo, P. and Li, J., Monosynaptic connections between neurons of trigeminal mesencephalic nucleus and jaw-closing motoneurons in the rat: an intracellular horseradish peroxidase labelling study, 267
Lynch, G.R., see Zlomanczuk, P., 94

Ma, S. and Long, J.P., Central noradrenergic activity is responsible for nitroglycerin-induced cardiovascular effects in the nucleus tractus solitarii, 297
Mackinnon, S.E., see Evans, P.J., 315
Maeda, N., see Matsuoka, N., 233
Makino, A.P., see Evans, P.J., 315
Maler, L., see Zupanc, G.K.H., 249
Manzanares, J., see Lookingland, K.J., 82
Margraf, R.R., see Zlomanczuk, P., 94
Martin, B.R., see Lichtman, A.H., 309
Masuyama, Y., see Ueyama, T., 293
Matsuoka, N., Maeda, N., Ohkubo, Y. and Yamaguchi, I., Differential effects of physostigmine and pilocarpine on the spatial memory deficits produced by two septo-hippocampal deafferentations in rats, 233
Mayer, A.D., see Cunningham, S.T., 337
McGeer, E.G., see Ousky, J.R., 322
McGuire, M.T., see Raleigh, M.J., 181
Metherate, R. and Ashe, J.H., Basal forebrain stimulation modifies auditory cortex responsiveness by an action at muscarinic receptors, 163
Mizuno, N., see Yasui, Y., 139
Mocchetti, I., see Reynolds, M.E., 149
Moore, K.E., see Lookingland, K.J., 82

Naegele, J.R. and Barnstable, C.J., A carbohydrate epitope defined by monoclonal antibody VC1.1 is found on

Brain Research, 559 (1991) 358–359
Elsevier

Author Index

Andrews, B.J., see Granat, M.H., 344
Aoki, E., Semba, R. and Kashiwamata, S., Evidence for the presence of L-arginine in the glial components of the peripheral nervous system, 159
Appenteng, K., see Saha, S., 145
Ashe, J.H., see Metherate, R., 163

Bain, J.R., see Evans, P.J., 315
Barnstable, C.J., see Naegele, J.R., 118
Batten, T.F.C., see Saha, S., 145
Baudry, M., see Tocco, G., 168
Baudry, M., see Tocco, G., 331
Baxendale, R.H., see Granat, M.H., 344
Bayer, V.E. and Pickel, V.M., GABA-labeled terminals form proportionally more synapses with dopaminergic neurons containing low densities of tyrosine hydroxylase-immunoreactivity in rat ventral tegmental area, 44
Berecek, K.H., see Swords, B.H., 10
Besharse, J.C., see Grace, M.S., 56
Besson, M.J., see Le Boulch, N.L., 200
Betancur, C., see Neveu, P.J., 347
Biella, G., Panara, C., Pecile, A. and Sotgiu, M.L., Facilitatory role of calcitonin gene-related peptide (CGRP) on excitation induced by substance P (SP) and noxious stimuli in rat spinal dorsal horn neurons. An iontophoretic study in vivo, 352
Blaustein, J.D., see Delville, Y., 191
Blomqvist, L., see Sjögren, A., 276
Boldry, R.C., Willins, D.L., Wallace, L.J. and Uretsky, N.J., The role of endogenous dopamine in the hypermotility response to intra-accumbens AMPA, 100
Boyes, B.E., see Ousky, J.R., 322
Brammer, G.L., see Raleigh, M.J., 181
Brunello, N., see Reynolds, M.E., 149

Cahill, G.M., see Grace, M.S., 56
Connelly, C.D., see Raffa, R.B., 211
Crain, S.M., see Shen, K.-F., 130
Creese, I., see Tepper, J.M., 283
Cunningham, S.T., Steinman, J.L., Whipple, B., Mayer, A.D. and Komisaruk, B.R., Differential roles of hypogastric and pelvic nerves in the analgesic and motoric effects of vaginocervical stimulation in rats, 337

DeGeorge, J.J., see Nariai, T., 1
Delville, Y. and Blaustein, J.D., A site for estradiol priming of progesterone-facilitated sexual receptivity in the ventrolateral hypothalamus of female guinea pigs, 191
Devgan, K.K., see Tocco, G., 331
Dragunow, M., see Jansen, K.L.R., 172
Duffy, P., see Samathanam, G.K., 37

Evans, P.J., Bain, J.R., Mackinnon, S.E., Makino, A.P. and Hunter, D.A., Selective reinnervation: a comparison of recovery following microsuture and conduit nerve repair, 315

Faull, R.L.M., see Jansen, K.L.R., 172

Gábriel, R., Zhu, B. and Straznicky, C., Tyrosine hydroxylase-immunoreactive elements in the distal retina of *Bufo marinus*: a light and electron microscopic study, 225
Gauchy, C., see Le Boulch, N.L., 200
Grace, M.S., Cahill, G.M. and Besharse, J.C., Melatonin deacetylation: retinal vertebrate class distribution and *Xenopus laevis* tissue distribution, 56
Granat, M.H., Nicol, D.J., Baxendale, R.H. and Andrews, B.J., Dishabituation of the flexion reflex in spinal cord-injured man and its applicatioin in the restoration of gait, 344
Grant, S.J. and Highfield, D.A., Extracellular characteristics of putative cholinergic neurons in the rat laterodorsal tegmental nucleus, 64
Gunnet, J.W., see Lookingland, K.J., 82

Hama, H., see Yamaguchi, K., 261
Hamada, M., see Ueyama, T., 293
Hano, T., see Ueyama, T., 293
Hauge, S.A., see Tocco, G., 331
Heiligenberg, W., see Zupanc, G.K.H., 249
Highfield, D.A., see Grant, S.J., 64
Ho, L.T., see Zochodne, D.W., 89
Hollingworth, R.M., see Raffa, R.B., 211
Hori, T., see Tanaka, T., 154
Hunter, .A., see Evans, P.J., 315

Ireland, L.M., see Lookingland, K.J., 82

Jansen, K.L.R., Faull, R.L.M., Dragunow, M. and Leslie, R.A., Autoradiographic distribution of sigma receptors in human neocortex, hippocampus, basal ganglia, cerebellum, pineal and pituitary glands, 172

Kalivas, P.W., see Samathanam, G.K., 37
Kalivas, P.W., see Sorg, B.A., 29
Karius, D.R., Ling, L. and Speck, D.F., Lesions of the rostral dorsolateral pons have no effect on afferent-evoked inhibition of inspiration, 22
Kashiwamata, S., see Aoki, E., 159
Kayahara, T., see Yasui, Y., 139
Kendig, J.J., see Woodley, S.J., 17
Komisaruk, .R., see Cunningham, S.T., 337
Kondo, S., see Tanaka, T., 154
Korn, H., see Petrov, T., 75
Ku, Y.-H. and Wang, X.-Q., Favored patterns in spontaneous spike trains, 241

Lamour, Y., see Nariai, T., 1
Le Boulch, N.L., Truong-Ngoc, N.A., Gauchy, C. and Besson, M.J., In vivo release of newly synthesized [^{3}H]GABA in the substantia nigra of the rat: relative contribution of GABA striato-pallidonigral afferents and nigral GABA neurons, 200
Le Moal, M., see Neveu, P.J., 347
Ledeen, R.W., see Shen, K.-F., 130
Leslie, R.A., see Jansen, K.L.R., 172
Li, J., see Luo, P., 267
Lichtman, A.H. and Martin, B.R., Cannabinoid-induced antinociception is mediated by a spinal α_2-noradrenergic mechanism, 309
Lindå, H., see Sjögren, A., 276
Ling, L., see Karius, D.R., 22
Long, J.P., see Ma, S., 297
Lookingland, K.J., Ireland, L.M., Gunnet, J.W., Manzanares, J., Tian, Y. and Moore, K.E., 3-Methoxy-4-hydroxyphenylethyleneglycol concentrations in discrete hypothalamic nuclei reflect the activity of noradrenergic neurons, 82
Lu, C.-L., see Shaikh, M.B., 109
Luo, P. and Li, J., Monosynaptic connections between neurons of trigeminal mesencephalic nucleus and jaw-closing motoneurons in the rat: an intracellular horseradish peroxidase labelling study, 267
Lynch, G.R., see Zlomanczuk, P., 94

Ma, S. and Long, J.P., Central noradrenergic activity is responsible for nitroglycerin-induced cardiovascular effects in the nucleus tractus solitarii, 297
Mackinnon, S.E., see Evans, P.J., 315
Maeda, N., see Matsuoka, N., 233
Makino, A.P., see Evans, P.J., 315
Maler, L., see Zupanc, G.K.H., 249
Manzanares, J., see Lookingland, K.J., 82
Margraf, R.R., see Zlomanczuk, P., 94
Martin, B.R., see Lichtman, A.H., 309
Masuyama, Y., see Ueyama, T., 293
Matsuoka, N., Maeda, N., Ohkubo, Y. and Yamaguchi, I., Differential effects of physostigmine and pilocarpine on the spatial memory deficits produced by two septo-hippocampal deafferentations in rats, 233
Mayer, A.D., see Cunningham, S.T., 337
McGeer, E.G., see Ousky, J.R., 322
McGuire, M.T., see Raleigh, M.J., 181
Metherate, R. and Ashe, J.H., Basal forebrain stimulation modifies auditory cortex responsiveness by an action at muscarinic receptors, 163
Mizuno, N., see Yasui, Y., 139
Mocchetti, I., see Reynolds, M.E., 149
Moore, K.E., see Lookingland, K.J., 82

Naegele, J.R. and Barnstable, C.J., A carbohydrate epitope defined by monoclonal antibody VC1.1 is found on

N-CAM and other cell adhesion molecules, 118
Nakano, K., see Yasui, Y., 139
Nariai, T., DeGeorge, J.J., Lamour, Y. and Rapoport, S.I., In vivo brain incorporation of [1-^{14}C]arachidonate in awake rats, with or without cholinergic stimulation, following unilateral lesioning of nucleus basalis magnocellularis 1
Neveu, P.J., Betancur, C., Vitiello, S. and Le Moal, M., Sex-dependent association between immune function and paw preference in two substrains of C3H mice, 347
Nicol, D.J., see Granat, M.H., 344
Nishio, I., see Ueyama, T., 293

Ohkubo, Y., see Matsuoka, N., 233
Ousky, J.R., Boyes, B.E., Walker, D.G. and McGeer, E.G., Cytomegalovirus infection of the developing brain alters catecholamine and indoleamine metabolism, 322
Orr, N., see Raffa, R.B., 211

Panara, C., see Biella, G., 352
Pecile, A., see Biella, G., 352
Petrov, T., Seitanidou, T., Triller, A. and Korn, H., Differential distribution of GABA- and serotonin-containing afferents on an identified central neuron, 75
Pickel, V.M., see Bayer, V.E., 44
Pollack, D.B., see Raleigh, M.J., 181

Raffa, R.B., Orr, N., Connelly, C.D. and Hollingworth, R.M., XAMI and DCDM, agonists at cAMP-associated octopamine receptors in cockroach nerve cord, produce centrally mediated antinociception in mice, 211
Raleigh, M.J., McGuire, M.T., Brammer, G.L., Pollack, D.B. and Yuwiler, A., Serotonergic mechanisms promote dominance acquisition in adult male vervet monkeys, 181
Rapoport, S.I., see Nariai, T., 1
Remahl, S., see Sjögren, A., 276
Reynolds, M.E., Brunello, N., Mocchetti, I. and Wrathall, J.R., Localization of nerve growth factor receptor mRNA in contused rat spinal cord by in situ hybridization, 149
Riekkinen, M., see Riekkinen Jr., P., 304
Riekkinen, P., see Riekkinen Jr., P., 304
Riekkinen Jr., P., Sirviö, J., Riekkinen, M. and Riekkinen, P., EEG changes induced by acute and chronic quisqualic or ibotenic acid nucleus basalis lesions are stabilized by tacridine 304
Risling, M., see Sjögren, A., 276

Saha, S., Appenteng, K. and Batten, T.F.C., Light and electron microscopical localisation of 5-HT-immunoreactive boutons in the rat trigeminal motor nucleus, 145
Samathanam, G.K., White, S.R., Kalivas, P.W. and Duffy, P., Effects of 5-hydroxytryptophan on extracellular serotonin in the spinal cord of rats with experimental allergic encephalomyelitis, 37
Schwartz, D.H., see Tepper, J.M., 283
Seitanidou, T., see Petrov, T., 75
Semba, R., see Aoki, E., 159
Shaikh, M.B., Lu, C.-L. and Siegel, A., An enkephalinergic mechanism involved in amygdaloid suppression of affective defence behavior elicited from the midbrain periaqueductal gray in the cat, 109
Shen, K.-F., Crain, S.M. and Ledeen, R.W., Brief treatment of sensory ganglion neurons with GM1 ganglioside enhances the efficacy of opioid excitatory effects on the action potential, 130
Shors, T.J., see Tocco, G., 168
Siegel, A., see Shaikh, M.B., 109
Sirviö, J., see Riekkinen Jr., P., 304
Sjögren, A., Thelestam, M., Blomqvist, L., Lindå, H., Remahl, S. and Risling, M., Extravasation of staphylococcal α-toxin in normal and injured CNS regions lacking blood–brain barrier function: observations after ventral root replantation, 276
Sorg, B.A. and Kalivas, P.W., Effects of cocaine and footshock stress on extracellular dopamine levels in the ventral striatum, 29
Sotgiu, M.L., see Biella, G., 352
Speck, D.F., see Karius, D.R., 22
Steinman, J.L., see Cunningham, S.T., 337
Straznicky, C., see Gábriel, R., 225
Swords, B.H., Wyss, J.M. and Berecek, K.H., Central vasopressin receptors are upregulated by deoxycorticosterone acetate, 10

Tanaka, S., see Tanaka, T., 154
Tanaka, T., Kondo, S., Hori, T., Tanaka, S. and Yonemasu, Y., Various hippocampal lesions induced by multi-fractional ibotenic acid injections and amygdala kindling in rats, 154
Tepper, J.M., Creese, I. and Schwartz, D.H., Stimulus-evoked changes in neostriatal dopamine levels in awake and anesthetized rats as measured by microdialysis, 283
Thelestam, M., see Sjögren, A., 276
Thompson, R.F., see Tocco, G., 168
Thompson, R.F., see Tocco, G., 331
Tian, Y., see Lookingland, K.J., 82
Tocco, G., Devgan, K.K., Hauge, S.A., Weiss, C., Baudry, M. and Thompson, R.F., Classical conditioning selectively increases AMPA receptor binding in rabbit hippocampus, 331
Tocco, G., Shors, T.J., Baudry, M. and Thompson, R.F., Selective increase of AMPA binding to the AMPA/quisqualate receptor in the hippocampus in response to acute stress, 168
Triller, A., see Petrov, T., 75
Truong-Ngoc, N.A., see Le Boulch, N.L., 200

Ueyama, T., Hano, T., Hamada, M., Nishio, I. and Masuyama, Y., New role of nerve growth factor — an inhibitory neuromodulator of adrenergic transmission, 293
Uretsky, N.J., see Boldry, R.C., 100

Vitiello, S., see Neveu, P.J., 347

Walker, D.G., see Ousky, J.R., 322
Wallace, L.J., see Boldry, R.C., 100
Wang, X.-Q., see Ku, Y.-H., 241
Weiss, C., see Tocco, G., 331
Whipple, B., see Cunningham, S.T., 337
White, S.R., see Samathanam, G.K., 37
Willins, D.L., see Boldry, R.C., 100
Woodley, S.J. and Kendig, J.J., Substance P and NMDA receptors mediate a slow nociceptive ventral root potential in neonatal rat spinal cord, 17
Wrathall, J.R., see Reynolds, M.E., 149
Wyss, J.M., see Swords, B.H., 10

Yamaguchi, I., see Matsuoka, N., 233
Yamaguchi, K. and Hama, H., Separation of periventricular dopaminergic and α-adrenergic systems from the vasopressin-secreting mechanisms activated by prostaglandin D_2, 261
Yasui, Y., Nakano, K., Kayahara, T. and Mizuno, N., Non-dopaminergic projections from the substantia nigra pars lateralis to the inferior colliculus in the rat, 139
Yonemasu, Y., see Tanaka, T., 154
Yuwiler, A., see Raleigh, M.J., 181

Zhu, B., see Gábriel, R., 225
Zlomanczuk, P., Margraf, R.R. and Lynch, G.R., In vitro electrical activity in the suprachiasmatic nucleus following splitting and masking of wheel-running behavior, 94
Zochodne, D.W. and Ho, L.T., Unique microvascular characteristics of the dorsal root ganglion in the rat, 89
Zupanc, G.K.H., Maler, L. and Heiligenberg, W., Somatostatin-like immunoreactivity in the region of the prepacemaker nucleus in weakly electric knifefish, *Eigenmannia:* a quantitative analysis, 249